

SUSY Searches at Tevatron Collider

C. Pagliarone

*University of Cassino & INFN of Pisa,
via di Biasio, 43 - 03043 Cassino, Italy
E-mail: pagliarone@fnal.gov*

This article presents recent results of searches for Supersymmetry using the CDF and the DØ detector at the Fermilab Tevatron Collider. Described are the Tevatron searches for third generation scalar quarks and for supersymmetric signatures involving photons. All the reported results have been obtained assuming theoretical models in which \mathcal{R} -parity is conserved.

1 Introduction

1.1 Supersymmetry

Although at present, the Standard Model (SM) provides a remarkably successful description of known phenomena, it seems to be, most likely, a low energy effective theory of spin-1/2 matter fermions interacting via spin-1 gauge bosons¹. An excellent candidate of a new theory, able to describe physics at arbitrarily high energies, is Supersymmetry (SUSY)². SUSY is a larger space-time symmetry, that relates bosons to fermions. Even if we don't have direct experimental evidences of SUSY, there are remarkable theoretical properties that provides ample motivation for its study. SUSY describe electroweak data equally well than SM but, in addition, allows the unification of the gauge couplings constants, the unification of the Yukawa couplings and do not require the incredible fine tuning, endemic to Higgs sector of the SM. Naturally, SUSY cannot be an exact symmetry of the nature, as none of the predicted spin 0 partners of the quarks or leptons and none of the spin 1/2 partners of the gauge bosons have been observed so far³.

In Supersymmetry fermions can couple to a sfermion and a fermion, violating lepton and/or baryon number. To avoid this problem, a new quantum number, the \mathcal{R} -parity, has been introduced⁴. For SM particles $\mathcal{R} = +1$, for the SUSY partners $\mathcal{R} = -1$. \mathcal{R} -parity conservation has deep phenomenological consequences⁵. SUSY particles can be only produced in pairs; the lightest supersymmetric particle (LSP) does exist and it is stable and interacts very weakly with the ordinary matter, leading to a robust missing transverse energy signature (\cancel{E}_T); the LSP is a natural candidate for the dark matter. In the present article, we present a review of recent Tevatron searches performed assuming \mathcal{R} -parity conservation.

1.2 Gauge Mediated SUSY Breaking Models

Theories with gauge-mediated supersymmetry breaking (GMSB) provide an interesting alternative scenario. In GMSB^{6,7}, the dynamical supersymmetry breaking (DSB) is mediated by gauge interactions. In recent years, many mechanism for DSB have been found and realistic models have been constructed. This class of models assumes that supersymmetry is broken with a scale \sqrt{F} in a sector of the theory




Figure 1: Jet multiplicity and E_T distribution after all cuts have been applied (CDF experiment).




Figure 2: 95% C.L. limit as function of $M_{\tilde{\chi}_1^0}$ and $M_{\tilde{t}_1}$, for direct $\tilde{t}_1\tilde{t}_1$ production, assuming $BR(\tilde{t}_1 \rightarrow c\tilde{\chi}_1^0) = 100\%$ (CDF experiment).

which contains heavy non-Standard-Model particles. This sector then couples to a set of particles with Standard Model interactions, called messengers, which have a mass of order M . The mass splitting, between the superpartners in the messenger multiplets, depends by \sqrt{F} and the SUSY particles get their masses via gauge interactions, so there are no flavor changing neutral currents. These theories have a very distinctive phenomenological features. The typical SUSY spectra is different from those in the SUGRA models; the LSP is the gravitino \tilde{G} (in SUGRA $M_{\tilde{G}} \sim 1$ TeV), the next lightest supersymmetric particle (NLSP) has a lifetime that can vary strongly from model to model ($1\mu < c\tau < \text{several Km}$) and decays into \tilde{G} .

2 Search for scalar top

Search for scalar top is particularly interesting since, in many SUSY models, the top-squark eigenstate \tilde{t}_1 (stop) is expected to be the lightest squark⁸. The strong Yukawa coupling, between top/stop and Higgs fields, gives rise, in fact, to potentially large mixing effects and mass splitting. The CDF experiment has recently performed two different searches for scalar top: direct $\tilde{t}_1\tilde{t}_1$ production, assuming a branching ratio $BR(\tilde{t}_1 \rightarrow c\tilde{\chi}_1^0) = 100\%$, and a search for supersymmetric decays of the top quark: $t \rightarrow \tilde{t}_1\tilde{\chi}_1^0$, with $\tilde{t}_1 \rightarrow b\tilde{\chi}_1^\pm$.

2.1 Search for direct Stop pair production

Whether \mathcal{R} -parity is conserved or not, at Tevatron, stop quarks are produced in pairs via gg and $q\bar{q}$ fusion. The LO diagonal pair production cross section depends mainly on stop mass and very little on other SUSY parameters such as gluino mass,




Figure 7: The E_T spectrum for diphoton events with $E_T^\gamma > 12 \text{ GeV}$ and with $E_T^\gamma > 25 \text{ GeV}$ in the data from CDF detector.




Figure 8: 95% C.L. excluded region (M_2 vs $\tan\beta$) for the Light Gravitino LSP scenario (CDF experiment).

tron, the sbottom can only decay into $b\tilde{\chi}_1^0$. The decays of the sbottom into $\tilde{t}_1 W$, with a virtual W , $\tilde{t}_1 \tilde{\chi}_1^-$, or into $\tilde{t}_1 H^\pm$ are kinematically suppressed. The signature for the process under study is therefore 2 b -quarks plus \cancel{E}_T . This final state is similar to those of two published $D\bar{O}$ analysis: the search for third generation leptoquark ($\nu\nu b\bar{b}$)¹⁵, and the search for direct scalar top production ($\tilde{t}_1 \tilde{t}_1 \rightarrow c\bar{c}\tilde{\chi}_1^0\tilde{\chi}_1^0$)¹⁴. The limits on sbottom production have been obtained combining these two samples. The number of events observed in the data is 5; 6.0 ± 1.3 events are expected from SM background. Figure 6 plots the 95% C.L. limit, as function of $M_{\tilde{b}_1}$ and $M_{\tilde{\chi}_1^0}$. A lower limit on the sbottom mass of $M_{\tilde{b}} > 115 \text{ GeV}/c^2$ which is valid for $M_{\tilde{\chi}_1^0} < 20 \text{ GeV}/c^2$ has been placed.

4 Photon enriched SUSY

In the last few years, triggered by the CDF $ee\gamma\gamma + \cancel{E}_T$ event candidate¹⁶, many models of new physics, predicting photon enriched final states have been introduced. CDF and $D\bar{O}$ have systematically searched for events containing two photons in the final state: $\gamma\gamma + X$, where X can be a jet, a b -tag, a lepton (e, μ, τ) or \cancel{E}_T .

4.1 Light Gravitino LSP

In the Minimal Gauge Mediated SUSY Breaking Model (MGM), as in SUGRA, the supersymmetry breaking occurs in a hidden sector. Unlike the SUGRA models, in which $\Lambda_{SUSY} \sim 10^{11}$, in MGM $\Lambda_{SUSY} \sim 10^5 - 10^9$, making the Gravitino the lightest supersymmetric particle ($M_{\tilde{G}} \sim 1 \div 10^2 \text{ eV}$). The Next-to-lightest SUSY




Figure 9: The \cancel{E}_T spectrum for diphoton events with $E_T^{\gamma}(1) > 20 \text{ GeV}$ and $E_T^{\gamma}(2) > 12 \text{ GeV}$ in the data from $D\phi$ experiment.




Figure 10: 95% C.L. excluded region (μ vs M_2) for the Light Gravitino LSP scenario ($D\phi$ detector); the dashed area shows the region suggested to explain the CDF $ee\gamma\gamma + \cancel{E}_T$ event.

particle (NLSP), which in the models is the neutralino ($\tilde{\chi}_1^0$), will decay dominantly via $\tilde{\chi}_1^0 \rightarrow \gamma \tilde{G}$. Therefore, any pair produced sparticle will produce two photons and \cancel{E}_T ¹⁷. In this model the $\tilde{\chi}_1^0$ lifetime depends on $M_{\tilde{G}}$ and is assumed to be less than 1 KeV. The gaugino masses are defined by M_2 , $\tan\beta$ and $\text{sign}\mu$ and $M_{\tilde{\chi}_1^\pm} \sim M_{\tilde{\chi}_1^0} \sim M_2$. In order to set a limit in this MGM model, the results of the counting experiment in the $\gamma\gamma \cancel{E}_T$ have been used. Events are asked to have 2 photons with $E_T^\gamma > 12 \text{ GeV}$ ($|\eta| < 1$) and $\cancel{E}_T > 35$ or $E_T^\gamma > 25 \text{ GeV}$ ($|\eta| < 1$) and $\cancel{E}_T > 25$. For $\cancel{E}_T > 35 \text{ GeV}$, 1 event passes the cuts, when the expected background is 0.5 ± 0.1 (see figure 7)¹⁶. Figure 8 shows the contour plot of the 95% C.L. excluded region as function of $\tan\beta$ and M_2 .

An analogous search has been performed by $D\phi$ Collaboration by selecting events with $E_T^{\gamma 1} > 20 \text{ GeV}$ and $E_T^{\gamma 2} > 12 \text{ GeV}$ ($|\eta| < 1.2$ or $1.5 < |\eta| < 2.0$). For $\cancel{E}_T > 25 \text{ GeV}$, 2 events are observed when 2.3 ± 0.9 are expected from SM background processes (see figure 9). Figure 10 plots the 95% C.L. excluded region (μ vs M_2).

4.2 Neutralino Radiative Decay

The $\gamma \cancel{E}_T + n$ jets event topology may also arise in MSSM, in some region of the parameter space, where the radiative decay $\tilde{\chi}_2^0 \rightarrow \gamma \tilde{\chi}_1^0$ dominates. Depending on the number of $\tilde{\chi}_2^0$, involved in the process, it will be possible to have one or more γ in the final state. $D\phi$ has searched for such signal using 99 pb^{-1} of data, assuming that slepton masses are heavy, $M_{\tilde{\chi}_2^0} - M_{\tilde{\chi}_1^0} > 20 \text{ GeV}/c^2$ and $\mathcal{BR}(\tilde{\chi}_2^0 \rightarrow \gamma \tilde{\chi}_1^0) = 100\%$. In order to select $\gamma \cancel{E}_T + \geq 2$ jets event candidates, we require to have at least one identified photon with $E_T^\gamma > 20 \text{ GeV}$, $\cancel{E}_T > 25 \text{ GeV}$ (see fig. 11) and two or more jets ($E_T^{\text{jet}} > 20 \text{ GeV}$). Then the event selection has been optimized in the \cancel{E}_T , H_T plane ($\cancel{E}_T > 45 \text{ GeV}$, $H_T > 220 \text{ GeV}$), resulting in 5 events passing the above cuts. 8.1 ± 5.8 events are expected from SM background processes. The 95% C.L. upper limit on $\sigma \times \mathcal{BR}$ as function of $M_{\tilde{q}}$ ($M_{\tilde{q}} = M_{\tilde{g}}$) is given in figure 12.




Figure 11: The E_T spectrum for events with one photon ($E_T^\gamma(1) > 20 \text{ GeV}$) and two or more jets ($E_T^{\text{jet}} > 20 \text{ GeV}$) in the data from $D0$ experiment.




Figure 12: The 95% C.L. upper limit on $\sigma \times BR$ as function of $M_{\tilde{q}} = m_{\tilde{q}}$. The hatched band represents the range of theoretical cross section for different sets of MSSM parameter values ($D0$ experiment).

4.3 The Higgsino LSP scenario

The present Higgsino LSP scenario assumes MSSM without sfermion/scalar masses unification. $\tilde{\chi}_1^0$ is an Higgsino-like LSP ($\tan\beta = 1.2$), and $\tilde{\chi}_2^0$ is photino-like ($M_1 = M_2$). So doing this model predicts a large branching ratio for the process $\tilde{\chi}_2^0 \rightarrow \gamma \tilde{\chi}_1^0$ and a light stop. A branching ratio of $\mathcal{BR}(t_1 \rightarrow c \tilde{\chi}_1^0) = 100\%$ is assumed. We end up with a topology containing $\gamma b + \cancel{E}_T$ in the final state: $p\bar{p} \rightarrow C_1 N_2 \rightarrow \tilde{t} b \gamma N_1 \rightarrow c b \gamma N_1 N_1$. CDF has searched for such events requiring a photon with $E_T^\gamma > 25 \text{ GeV}$, a SVX b -tag and $\cancel{E}_T > 25 \text{ GeV}$. To increase the sensitivity we then required the photon to be not opposite to the \cancel{E}_T , and $\cancel{E}_T > 40 \text{ GeV}$ (see figure 13). We see 2 events; this allows to rule out more than approximately 7 events of anomalous production. Figure 14 shows the limit, plotted as function of gluino mass. In figure 15 we present the limit on direct production of $\tilde{\chi}_2^\pm \tilde{\chi}_2^0$.

4.4 Gauge-mediated SUSY with $\gamma b + \cancel{E}_T$

Another way to produce the $\gamma b + \cancel{E}_T$ signature comes from a GMSB model. In this model, the gravitino is light and becomes the LSP. The $\tilde{\chi}_1^0$ is a higgsino and may decay into either a gravitino and a photon ($\tilde{\chi}_1^0 \rightarrow \tilde{G} \gamma$) or into a gravitino and an Higgs boson ($\tilde{\chi}_1^0 \rightarrow \tilde{G} h$), with the Higgs decaying into $b\bar{b}$. Since we have two $\tilde{\chi}_1^0$ in each event this will give rise to a $\gamma b + \cancel{E}_T$ signature. The CDF limits obtained in this scenario are shown in figure 16.

5 Conclusions

Extensive searches of supersymmetric signals have been done at Tevatron Collider. No positive results have been found so far showing that the data are consistent with




Figure 13: The E_T distribution in the data from CDF detector, compared to the SUSY Higgsino LSP baseline model.




Figure 14: The limits on the $\sigma \times \mathcal{BR}$ for SUSY production in the Higgsino LSP model (CDF experiment).

the SM expectation. Further regions of SUSY parameter space, assuming different SUSY models, have been excluded. With the Run II¹⁸ upgrades, providing an higher acceptance and higher luminosity, it will be possible to probe larger region of SUSY parameter space¹⁹.

References

1. G. Altarelli, CERN-TH-98-348, **hep-ph/9811456** (1998).
2. S.P. Martin, **hep-ph/9709356**.
3. W. de Boer, **hep-ph/9808448**
4. H. Dreiner, *Pramana* **51**, 123 (1998).
5. J. Ellis *et al.*, *Phys. Lett. B* **150**, 142 (1985).
F. Vissani, IC-96-32, **hep-ph/9602395** (1995).
6. M. Dine *et al.*, *Nucl. Phys. Proc. Suppl.* **62**, 266-275 (1998).
7. G.F. Giudice, R. Rattazzi, CERN-TH-97-380, **hep-ph/9801271** (1998).
C. Kolda, *Nucl. Phys. Proc. Suppl.* **62**, 485 (1998).
8. Ken-ichi Hikasa, Makoto Kobayashi, *Phys. Rev. D* **36**, 724 (1987).
9. W. Beenakker *et al.*, *Nucl. Phys. B* **515**, 3 (1997).
10. S. Abachi *et al.* (DØ Collaboration), *Phys. Rev. Lett.* **76**, 2222 (1996).
11. J. Sender, *Phys. Rev. D* **54**, 3271 (1996).
J. D. Wells and G. L. Kane, *Phys. Rev. Lett.* **76**, 869 (1996).
G. Mahlon and G. L. Kane, *Phys. Rev. D* **55**, 2779 (1997).
M. Hosch *et al.*, *Phys. Rev. D* **58**, 034002 (1999).




Figure 15: The limits on the $\tilde{\chi}_2^\pm \tilde{\chi}_2^0$ cross section in the SUSY Higgsino LSP model (CDF experiment).




Figure 16: The limits on the SUSY $\sigma \times \mathcal{BR}$ in the MGMSB model (CDF experiment).

12. F. Abe *et al.* (CDF Collaboration), *Phys. Rev. D* **51**, 4623 (1995).
F. Abe *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **74**, 2626 (1995).
13. A. Bartl *et al.*, *Z. Phys.* **C76**, 549 (1997).
14. F. Abachi *et al.* (DØ Collaboration), *Phys. Rev. Lett.* **76**, 2222 (1996).
15. F. Abachi *et al.* (DØ Collaboration), *Phys. Rev. Lett.* **81**, 38 (1998).
16. F. Abe *et al.*, (CDF Collaboration), *Phys. Rev. D* **81**, 1971 (1998).
F. Abe *et al.*, (CDF Collaboration), [hep-ex/9806034](#).
17. S. Ambrosanio *et al.*, *Phys. Rev. D* **54**, 5395 (1996).
18. R. Blair *et al.*, (CDF-II Collaboration), FERMILAB-PUB-96-390-E (1996).
19. D. Amidei *et al.*, (TeV-2000 Study Group Collaboration),